

# Three new species of the genus *Argyresthia* Hübner, [1825] from Guatemala, with notes on host plant evolution and Nearctic taxa (Lepidoptera: Argyresthiidae)

Jacob A. Gorneau, Loren D. Jones, José Monzón-Sierra & Jason J. Dombroskie

## Abstract

Neotropical species of the genus *Argyresthia* Hübner, [1825] represent a reservoir of undescribed biodiversity and are the most poorly known members of this cosmopolitan genus. From a series of Guatemalan material, three new species are described: *Argyresthia quetzaltenangonella* Gorneau & Dombroskie, sp. nov., *Argyresthia guatemala* Gorneau & Dombroskie, sp. nov., *Argyresthia iridescentia* Gorneau & Dombroskie, sp. nov., representing the first Neotropical species to be described in over a century, and the first from Central America. A phylogeny is inferred using the barcode gene cytochrome c oxidase subunit I (COI), representing an initial foray into the internal relationships of the family. For species with known host plants and available molecular data, a stochastic character mapping analysis was conducted to estimate character states at internal nodes. Furthermore, this phylogeny allows for preliminary insights into the monophyly of the subgenus *Blastotere* Ratzeburg 1840. Systematic work on *Argyresthia* remains particularly salient as species in the genus run the gamut from being pests to being considered extinct. Furthermore, notes are provided regarding the status of select Nearctic species, such as the ongoing confusion in North America between *Argyresthia goedartella* (Linnaeus, 1758) and *A. calliphanes* Meyrick 1913. Focusing taxonomic efforts on *Argyresthia* of the Neotropics will lay the foundation for future work regarding biodiversity, ecology, and biogeography of the family.

**Keywords:** Lepidoptera, Argyresthiidae, new species, host plants, phylogenetics, Guatemala.

## Tres nuevas especies del género *Argyresthia* Hübner, [1825] de Guatemala, con notas sobre la evolución de la planta nutricia y taxas Neárticas (Lepidoptera: Argyresthiidae)

## Resumen

Las especies neotropicales del género *Argyresthia* Hübner, [1825] (Lepidoptera: Argyresthiidae) representan un reservorio de biodiversidad no descrita y son los miembros menos conocidos de este género cosmopolita. A partir de una serie de material guatemalteco se describen tres nuevas especies: *Argyresthia quetzaltenangonella* Gorneau & Dombroskie, sp. nov., *Argyresthia guatemala* Gorneau & Dombroskie, sp. nov., *Argyresthia iridescentia* Gorneau & Dombroskie, sp. nov., representando la primera especie neotropical descrita en más de un siglo y la primera de América Central. Se infiere una filogenia utilizando el gen del código de barras citocromo c oxidasa subunidad I (COI), lo que representa una incursión inicial en las relaciones internas de la familia. Para especies con plantas nutricias conocidas y datos moleculares disponibles, se realizó un análisis de mapeo de caracteres estocásticos para estimar los estados de caracteres en los nodos internos. Además, esta filogenia permite obtener información preliminar sobre la monofilia del subgénero *Blastotere* Ratzeburg 1840. El trabajo sistemático sobre *Argyresthia*

sigue siendo particularmente destacado ya que las especies del género van desde ser plagas hasta considerarse extintas. Además, se proporcionan notas sobre el estado de especies neárticas seleccionadas, como la confusión actual en América del Norte entre *Argyresthia goedartella* (Linnaeus, 1758) y *A. calliphanes* Meyrick 1913. Centrar los esfuerzos taxonómicos en *Argyresthia* del Neotrópico sentará las bases para el trabajo futuro con respecto a la biodiversidad, ecología y biogeografía de la familia.

**Palabras clave:** Lepidoptera, Argyresthiidae, nuevas especies, plantas nutricias, filogenética, Guatemala.

## Introduction

The enigmatic genus *Argyresthia* Hübner, [1825] belongs to the superfamily Yponomeutoidea and is the sole lineage of the family Arguresathiidae, totaling just over 200 described species (Lewis & Sohn, 2015; Liu et al. 2017). While small in size, species of *Argyresthia* exhibit a variety of forewing patterns ranging from high-contrast metallics to dull silvers, to mottled browns, and are sometimes colloquially referred to as “shiny head-standing moths” for their resting posture. Currently, most species are Palearctic in distribution (Lewis & Sohn, 2015; Liu et al. 2017). Although it has been suggested by Heppner (2008) that the total diversity of the family may “exceed 450 species”, the recent description of forty-three new species from China alone indicates this is likely to be a lower estimate (Liu et al. 2017). The subgenus *Blastotere* Ratzeburg, 1840 was more formally characterized by Liu et al. (2017) and represents the sole subgenus aside from the nominate subgenus. It contains twenty-five species (Table I), in accordance with Bengtsson & Johansson (2011) and Liu et al. (2017).

**Table I.** Species currently described formally in subgenus *Blastotere* Ratzeburg 1840.

Species	Authority
<i>Argyresthia (Blastotere) affinicineretra</i>	Liu, Wang & Li, 2017
<i>A. (B.) albaureola</i>	Liu, Wang & Li, 2017
<i>A. (B.) amiantella</i>	(Zeller, 1847)
<i>A. (B.) anthocephala</i>	Meyrick, 1936
<i>A. (B.) arceuthina</i>	Zeller, 1839
<i>A. (B.) aureola</i>	Liu, Wang & Li, 2017
<i>A. (B.) bergiella</i>	(Ratzeburg, 1840)
<i>A. (B.) chalcocausta</i>	Meyrick, 1936
<i>A. (B.) cineretra</i>	Liu, Wang & Li, 2017
<i>A. (B.) densa</i>	Liu, Wang & Li, 2017
<i>A. (B.) dilectella</i>	Zeller, 1847
<i>A. (B.) dolichocoremata</i>	Liu, Wang & Li, 2017
<i>A. (B.) glabratella</i>	(Zeller, 1847)
<i>A. (B.) idiograpta</i>	Meyrick, 1935
<i>A. (B.) illuminatella</i>	Zeller, 1839
<i>A. (B.) kulfani</i>	Bengtsson & Johansson, 2011
<i>A. (B.) laevigatella</i>	Herrich-Schäffer, 1855
<i>A. (B.) longipenella</i>	Liu, Wang & Li, 2017
<i>A. (B.) ornatipenella</i>	Moriuti, 1974
<i>A. (B.) praecocella</i>	Zeller, 1839
<i>A. (B.) punctireticulata</i>	Liu, Wang & Li, 2017
<i>A. (B.) sporadolepis</i>	Liu, Wang & Li, 2017
<i>A. (B.) svenssoni</i>	Bengtsson & Johansson, 2011
<i>A. (B.) trifasciata</i>	Staudinger, 1871

As larvae, species in the genus *Argyresthia* are miners and borers on gymnosperms and angiosperms. Host plant records currently exist for approximately one-third of the species in the genus

(Friese, 1969; Bengtsson & Johansson, 2011; Lewis & Sohn, 2015). Of these sixty-one species, forty-two of them feed on gymnosperms, and sixteen on Rosaceae (Friese, 1969; Bengtsson & Johansson, 2011; Lewis & Sohn, 2015). Three are cited as polyphagous (Lewis & Sohn, 2015). Liu et al. (2017) have suggested that the evolution of this group in China may be closely related to the diversity of gymnosperms in the region, noting the geographic similarities with areas of high *Argyresthia* diversity to high gymnosperm diversity, but this has yet to be evaluated in a phylogenetic context.

Knowledge of the systematics of this family is important to help understand pest management, climate change, and conservation. Some species are invasive pests of ornamental plants and fruit crops. *Argyresthia thuiella* (Packard, 1871) has been introduced from North America to Europe multiple times, where it has wreaked havoc on ornamental cupressaceous plants, such as that of *Thuja occidentalis* Linnaeus (Csóka, 2001). *Argyresthia trifasciata*, though native to Europe, appears to be spreading and has established itself as a pest in Hungary (Tiborne & Kálmán, 2000). Other species, such as *A. conjugella* Zeller, 1839, and *A. pruniella* (Clerck, 1759) are demonstrated pests of rosaceous fruit crops (Lampa, 2005; Shreyner, 2006; Schøyen, 1913; Ahlberg, 1927; Stapley, 1934; Belosel's Kaya, 1963; Korchagin, 1988; Sharma et al. 1988; Wimshurst, 1928; Alford, 1978; Carter, 1984; Alford, 2007; Jaastad, 2007; Loone et al. 2017). Climate change may have resulted in intensified outbreaks of *A. retinella* Zeller, 1839, a native birch (*Betula* spp.) feeder in northwestern Norway that until the 1990's was not known to exhibit an outbreak (Tenow et al. 1999). Conversely, one species, *A. castaneella* Busck, 1915, whose recorded host plant is the critically endangered American Chestnut (*Castanea dentata* (Marsh.) Borkh.), is now presumed extinct (World Conservation Monitoring Centre, 1996; Stritch, 2018). The relevance of this group to agricultural, horticultural, and natural systems makes applied and basic work on the Argylethiidae integral.

While the genus *Argyresthia* is well-studied in other regions of the world, particularly in Europe and Asia, a comprehensive review of the group in the Americas has not been completed in over one-hundred years, necessitating the taxonomic study of this group in the New World (Zeller, 1877; Bisck, 1907). Only seven species have been described from the Neotropics (Lewis & Sohn, 2015): *A. biruptella* Zeller, 1877, *A. carcinomatella* Zeller, 1877, *A. diffractella* Zeller, 1877, *A. ochridorsis* Zeller, 1877, and *A. percussella* Zeller, 1877 from Bogotá, Colombia; *A. conspersa* Butler, 1883 from Cauquenes, Chile; and *A. melिताula* Meyrick, 1918 described from La Cumbre, Colombia. Incidentally, no Central American *Argyresthia* species have been described or noted.

The goal of this work is to describe the Guatemalan holdings of *Argyresthia* in the Cornell University Insect Collection (CUIC) and to infer a phylogeny of the group based on the barcode gene, cytochrome oxidase c subunit I (COI). From this phylogeny, stochastic character mapping will be used to provide some preliminary insights into the evolution of host plant associations in the family. This study will also evaluate evidence for phylogenetic support of the subgenus *Blastotere* based on the COI inference. Lastly, some notes on the taxonomy of Nearctic species will be provided. Due to the estimated magnitude of unexplored diversity in this family and its worldwide distribution, this work also serves as a call-to-action for more to study the systematics of this family.

## Materials and Methods

Specimens were collected by JJD, Tim McCabe, and JMS in early 2014. Specimens and genitalia were photographed using Canon EOS 6D with a MacroPod Pro photo-stacking setup and images were automatically stacked using ZereneStacker. Photos were post-processed using Adobe Photoshop.

Dissections were made by removing the abdomen with fine forceps and placing it in a 1.5 mL plastic tube with approximately 1 mL of 10% potassium hydroxide (KOH) solution. The tube was then placed in a heat block at 81°C and was checked at twenty-minute intervals until the abdomen was translucent. The abdomen was then removed from the heat and placed in a Syracuse watch glass with 70% ethanol. A small brush was used to remove scales that were not digested by KOH. Once the abdomen was brushed clean of scales, the specimen was soaked in chlorazol black for one full minute,

and then placed in eosin Y, microwaved for four seconds, and then left to sit in the microwaved eosin y dye for at least 10 minutes afterwards to ensure adequate setting of the dye. Terminology used for descriptions is in accordance with Liu et al. (2017), which largely follows Moriuti (1977).

For data not generated by this study, 130 sequences of the DNA barcoding region, COI, were downloaded from GenBank and the Barcode of Life Database (BOLD, Table S1). Single legs from a total of seven specimens were sent to BOLD (Centre for Biodiversity Genomics, Guelph, ON, CA) for sequencing of COI, and one additional sample was sequenced through the Cornell Genomics Facility (Ithaca, NY, USA). All sequences were submitted to GenBank through the BOLD interface. For all samples, the standard HCO2198/LCO1490 primer set was used (Folmer et al. 1994). Sequence accession information for those containing data generated by this study are available in Table II.

**Table II.** Sequence information generated in this study with accession numbers.

Genus	Species	JD Accession	GenBank Accession	BOLD Voucher
<i>Argyresthia</i>	<i>quetzaltenangonella</i> sp. nov.	JD23247	OM158441	CUICA018-19
<i>Argyresthia</i>	<i>guatemala</i> sp. nov.	JD13161	OM158438	CUICA010-19
<i>Argyresthia</i>	<i>guatemala</i> sp. nov.	JD17039	OM158444	CUICA014-19
<i>Argyresthia</i>	<i>guatemala</i> sp. nov.	JD17808	OM158442	CUICA016-19
<i>Argyresthia</i>	<i>guatemala</i> sp. nov.	JD16994	OM158437	CUICA012-19
<i>Argyresthia</i>	<i>guatemala</i> sp. nov.	JD17474	N/A	CUICA015-19
<i>Argyresthia</i>	<i>iridescentia</i> sp. nov.	JD22454	OM158440	CUICA017-19
<i>Argyresthia</i>	sp. nov.	JD13196	OM158439	CUICA011-19
<i>Argyresthia</i>	sp. nov.	JD17038	OM158443	CUICA013-19

Sequences were then aligned using the program MAFFT using the L-INS-i method (Katoh & Standley, 2013). Sequences were then manually examined and alterations to the MAFFT alignment were made as necessary in the program Mesquite (version 6.0) to ensure no stop codons were present in the alignment (Maddison & Maddison, 2018). A maximum likelihood phylogeny was inferred using IQ-TREE, a maximum likelihood phylogeny was inferred with 10,000 replicates and UFBoot values as nodal supports (Nguyen et al. 2015; Kalyaanamoorthy et al. 2017; Hoang et al. 2018). Using another method to examine convergence, a NEXUS block partitioned by codon position was exported from Mesquite for use in MrBayes. The analysis ran for 100,000,000 generations, with a burn in of 25%, following an inverse gamma distribution, with four chains. The phytools function cophylo was used to examine similarity among inferences.

Stochastic character mapping of host plants was implemented in RStudio (2022) using the packages phytools, corHMM, and geiger (Revell 2012; Pennell et al. 2014; Boyko & Beaulieu, 2021). The phylogeny was trimmed to species for which host plant data were available. While it is customary to include only one representative for each species in comparative phylogenetic analyses, multiple representatives were included for paraphyletic species. The tree was converted to an ultrametric topology to eliminate the bias of branch lengths on the inference. Model selection for host plant stochastic character mapping was conducted with the fitMk function for the equal rates, symmetric rates, and all rates different models. The equal rates model was selected via Akaike information criterion (AIC) values. Stochastic map trees were generated under the equal rates model using the function make.simmap for 1000 generations using a Markov Chain Monte Carlo approach with sampling every 10 generations, for a total of 100 stochastic map trees. From these 100 trees, posterior probabilities were calculated for all nodes. While there is likely less plasticity necessary for a host plant change from gymnosperm to gymnosperm versus a change between gymnosperm and angiosperm host plants, to not over parameterize the dataset, which already has limited resolution due to the single locus nature of this study, no further assumptions were made beyond that of the equal rates model. The character matrix, with corresponding character states is available in Table SII. Code available from <https://doi.org/10.5281/zenodo.6865177>.

## Results

### *Argyresthia quetzaltenangonella*, Gorneau & Dombroskie, sp. nov. (Figure 1)

Type material: Holotype, ♂, GUATEMALA: Quetzaltenango, Fuentes Georginas, 14.750, -91.480; 26-II-2014; J. J. Dombroskie, T. McCabe & J. Monzón leg.; MV/UV light: 2439 m; JD23247 (CUIC).

Description: Male adult (Figure 1A): Head with vertex white and roughly scaled. Frons cream-colored and smoothly scaled, becoming slightly whiter toward the proboscis. Labial palpus with first segment cream-colored, second segment distinctly whiter, and held freely, not erect in an upright position. Antenna with white eyecap obscuring approximately one-fourth of the eye; flagellum ground color white, striped with alternating dark brown scales. Thorax white, tegula golden orange. Forewing length 6.5 mm (n = 1). Forewing ground color just slightly off-white, with first two-thirds of wing from costa toward inner margin lightly reticulated with slightly off-white squares outlined in brownish-gold and interrupted basally by a thick golden orange band parallel to inner margin and costa which tapers distally about one-third of the way toward apex of wing; towards inner margin, ground color slightly whiter than elsewhere and interrupted at about same place where the golden orange band tapers by a broken light brown fascia which penetrates slightly into reticulated region of forewing, last one-fifth of wing near apex mottled golden brown with a series of dark scales marking apex before fringe, which are all dark grayish-brown. Hindwing length 5.0 mm (n = 1). Hindwing gray, becoming darker toward apex and inner margin a grayish-yellow color. Foreleg entirely dark brown ventrally, banded dorsally, predominantly white in ground color with dark brown bands occurring at the joints; midleg banded dorsally, as in foreleg, with this pattern slightly fainter ventrally; hindleg as in midleg. Abdomen pearly and with light grayish-white ground color. Coremata present and occupying approximately two-thirds of the abdomen in length (Figure 1B).

Male genitalia (Figure 1C-D): Socius covered with approximately 20 elongate scale-like setae, with three or four additional elongate setae on inner side of socius (Figure 1C). Tube analis tapering and subequal to width of valva. Valva rotund, with very few hairs. Phallus (Figure 1D) kinked basally, with cornutus visible in apical third; cornutus with seven small notches.

Female: Unknown.

Diagnosis: Similar to *Argyresthia biruptella* Zeller, 1877, described from Bogotá, Colombia, but with less maculation throughout the forewing, a faded reticulated pattern on the first two-thirds of the wing from the costa, and the presence of a golden orange band which runs parallel to the inner margin and costa extending about one-third of the way down the wing.

Distribution: One male specimen known from a cool high elevation forest in Quetzaltenango, Guatemala (2439 m).

Biology: See further discussion under Stochastic Character Mapping, but likely gymnosperm-feeding, as the ancestral state for the clade this species is inferred to belong to is Cupressaceae-feeding. *Cupressus lusitanica* Miller is present in the vicinity of this location.

Etymology: It is so named for the Quetzaltenango Department in Guatemala, where the type was found.

### *Argyresthia guatemala* Gorneau & Dombroskie, sp. nov. (Figure 2)

Type material: Holotype ♀, GUATEMALA, Alta Verapaz, Posada del Guardabarranco 15.300, -90.317; 02-III-2014; J. J. Dombroskie, T. McCabe & J. Monzón leg.; MV/UV light: 1549 m; JD13161 (CUIC).

Paratypes (2 ♂♂, 2 ♀♀), GUATEMALA, Alta Verapaz, 1 ♂ same data as holotype but JD17808 (Universidad del Valle de Guatemala Collection of Arthropods - UVGC). Suchitupéquez, Atitlan Reserve 14.548, -91.194, 1 ♂, 2 ♀♀, 23-II-2014, J. J. Dombroskie, T. McCabe & J. Monzón leg.; MV/UV light: 1561 m; JD16994 (CUIC), JD17039 (CUIC); JD17474 (Smithsonian National Museum of Natural History - NMNH).

Description: Female adult (Figure 2A): Head with vertex white and roughly scaled. Frons white to light beige and smoothly scaled. Labial palpus uniformly beige, sometimes darker than that of the frons, and held freely, not in an upright position. Antenna with white eyecap obscuring approximately one-fourth of the eye; flagellum ground color white, striped with alternating dark brown scales. Thorax white, tegula a light purplish brown. Forewing length 4.0 - 4.3 mm (n = 3). Forewing ground color cream colored, forewing from costa a mottled brown which becomes lightly reticulated toward the inner margin. About halfway down the inner margin, a small dark brown patch is present, with a smaller patch just opposite this on the costa. There are two more patches between the others, one located apically relative to the midpoint between the first two patches, and one located basally relative to the midpoint between the first two patches. Another patch about one-sixth of the way from the base of the wing that may just appear as a darker reticulation near the inner margin and another patch just slightly apical to that in the about halfway between the inner margin and the costa. Another dark patch on the apex of the wing. Hindwing length 3.3 - 3.5 mm (n = 3). Hindwing a uniform silver. Foreleg entirely dark brown ventrally, banded dorsally, predominantly white in ground color with dark brown bands occurring at the joints. Midleg banded dorsally, as in foreleg, with this pattern slightly fainter ventrally; hindleg as in midleg. Abdomen uniform silver as in hindwing.

Female genitalia (Figure 2B-C): Ductus bursae elongate, roughly 2.3 times the length of corpus bursae, and widening anteriorly approximately halfway to the corpus bursae (Figure 2B). Corpus bursae (Figure 2B) ovate and tapered distally as it nears the ductus bursae, with few maculations. Basal plate of signum ovoid and denticulate, with strongly sclerotized horn on either side of this plate slightly curved posteriorly. Horn of signum serrated regularly. Anterior to the horn of the signum, the basal plate remains ovoid but smaller in surface area, covering approximately one-third of the area of the basal posterior to the horn. Antrum and posterior apophyses visible in dissection image but damaged and indistinct image as ductus bursae separated from last abdominal segment in dissection process (Figure 2C).

Male adult (Paratypes): As in female, but with forewing length 3.5 mm and hindwing length 3.0 mm (n = 2). Genitalia damaged and unavailable for dissection.

Diagnosis: Looks most similar to *Argyresthia deletella* Zeller, 1873 (Figure 3A-C), described from Texas, USA, but without the brown fascia occasionally present in *A. deletella*, and a more overall mottled appearance. Ground color is overall much whiter than that of *Argyresthia deletella*, which has an overall browner ground color with whitish regions just nearer the inner margin. The signum is strongly serrated in *A. guatemala* but smooth to lightly serrated in *A. deletella*. Furthermore, while both species have a lightly sclerotized plate that is medially located on the signum, the signum in *A. guatemala* is larger and bisected by this structure, while the signum in *A. deletella* remains strongly sclerotized and is not bisected by this structure (Figure 3C).

Distribution: Three specimens (two female, one male) from Atitlan Reserve in Suchitupéquez, Guatemala, and two specimens (one female, one male) from Posada Del Guardabarranco, Alta Verapaz, Guatemala. Both locations are moderate in elevation, from 1549 - 1561 m.

Biology: See further discussion under Stochastic Character Mapping, but host plant ambiguous based on phylogenetic placement and known host plants for related species.

Etymology: It is so named for the country in which this species has been recorded.

#### *Argyresthia iridescentia* Gorneau & Dombroskie, sp. nov. (Figure 4)

Type material: Holotype, ♂, GUATEMALA: Quetzaltenango, Fuentes Georginas 14.750 -91.480; 26-II-2014; J. J. Dombroskie, T. McCabe & J. Monzón leg.; MV/UV light: 2439 m; JD22454 (CUIC).

Description: Male adult (Figure 4A): Head with vertex grayish white and roughly scaled. Frons uniformly light gold and smoothly scaled. Labial palpus uniformly light gold and held freely, not in an upright position. Antenna with white eyecap obscuring approximately one-fourth of the eye. Flagellum grayish white and striped with alternating gold scales. Thorax and tegula gold. Forewing length approximately 3.5 mm. Forewing ground color metallic gold at basal fifth of the wing and fading in

color apically, to match the pale gray-gold of hindwing. Gold ground color of wings can appear as a plumbeous brown when examined at certain angles or photographed. Four broken fasciae a darker metallic gold in coloration. First fascia appears one-fifth from the base of the wing at the inner margin and extends about halfway to the costa. Second fascia also originates from the inner margin, about halfway from the base of the wing and splits into two branches about one-third of the way from the inner margin toward the costa, as to create a V-shape. This fascia fades into the brownish-lead costa about three-fourths of the way from the inner margin. Third fascia originates at the inner margin where the fringe of scales on the forewing also begins and extends diagonally toward the apex and three-fourths of the way to the costa. Fourth fascia originates from the costa about halfway to the inner margin and appears much like a triangle. All fasciae may become worn, and appear as a few distinct spots, as in the left wing of the holotype. The apex of the wing has a small patch of five to ten very dark brown scales on the edge of the inner margin. These scales can be reduced to just a few scales from wear. Hindwing length 3.0 mm. Hindwing uniformly light gray-gold. Foreleg entirely dark brown ventrally, predominantly cream colored with slightly darkened bands at the joints dorsally. Midleg banded dorsally, as in foreleg, with this pattern slightly fainter ventrally. Hindleg as in midleg. Abdomen a pearly grayish white. Coremata not present.

Male genitalia (Figure 4B-C): socius covered with elongate scale-like setae (Figure 4B). Tuba analis stout, tapering and subequal to width of valva, valva rotund, and apparently hairless. Phallus kinked basally, with apical third of phallus containing cornutus (Figure 4C). Cornutus with seven small notches.

Female: Unknown.

Diagnosis: Looks most similar to *Argyresthia flexilis* Freeman, 1960, especially in that the wings can appear dull brown at one angle, and near gold in another. The general ground color of the wings is more plumbeous than that of *A. flexilis* and has noticeable maculations and fasciae, where *A. flexilis* is more uniformly patterned. *Argyresthia iridescentia* also has a patch of distinctly dark brown scales at the apex of the forewing.

Distribution: One male specimen known from a cool high elevation forest in Quetzaltenango, Guatemala (2439 m).

Biology: See further discussion under Stochastic Character Mapping, but likely gymnosperm-feeding, as the ancestral state for the clade this species is inferred to belong to is Pinaceae-feeding. *Abies guatemalensis* Rehder and several *Pinus* species can be found in the vicinity of this site.

Etymology: It is so named for the iridescent appearance of the wings which make them appear a pale lead brown at some angles, but golden at others.

## Phylogenetic Inference

A monophyletic *Argyresthiidae* with convergent topologies was supported in both the maximum likelihood and Bayesian inferences (Figure 6, UFBoot = 99; Figure S1, posterior probability = 1.00). While higher-level relationships were not consistent between the inferences, most species-level groupings were (Figure S2). The new species described in this paper were all recovered as distinct (in the case of *Argyresthia quetzaltenangonella* and *A. iridescentia*, with single representatives) or monophyletic where there were multiple representatives. Eleven out of twenty-five representatives were included from the subgenus *Blastotere*, which was recovered as paraphyletic. *Argyresthia goedartella* (Linnaeus 1758) and *A. calliphanes* Meyrick, 1913 formed a monophyletic group but were rendered paraphyletic in the groups examined here. Despite this, two representatives from Europe identified as *Argyresthia goedartella* formed a monophyletic group, Western North American representatives identified as *A. calliphanes* formed a monophyletic group, and Eastern North American representatives of *A. goedartella* and *A. calliphanes* formed a monophyletic group. *Argyresthia cupressella* Walsingham 1890, *Argyresthia trifasciae* Braun, 1910, *A. abies* Freeman, 1972, *A. glabratella*, *A. illuminatella*, *A. quadristrigella* Zeller, 1873 were all rendered paraphyletic in both inferences.

## Stochastic Character Mapping

The results of the stochastic character mapping, while inconclusive suggest a potentially Betulaceae-feeding origin for the family (Figure 7). There appear to be two predominant lineages within the Argyresthiidae, one ancestrally Rosaceae-feeding and one ancestrally Cupressaceae-feeding. Within the latter, there are two lineages ancestrally Pinaceae-feeding and one apparent evolution of angiosperm feeding in *Argyresthia glaucinella* Zeller, 1839. Sister to the clade ancestrally Cupressaceae-feeding *Argyresthia* is a grade of Betulaceae-feeding *Argyresthia*.

The results of the stochastic character mapping allow for some broad inferences about host plant use for species that do not currently have a recorded host plant (Figure 7). Species lacking a recorded host plant which fall into the gymnosperm-feeding clade include: *Argyresthia ruidosa* Braun, 1940, *A. austerella* Zeller, 1873, and *A. deletella* Zeller, 1873. Species which fall into the angiosperm-feeding clade include *A. atlanticella* Rebel, 1940, *A. notoleuca* (Turner, 1913), *A. assimilis* Moriuti, 1977, *A. pulchella* Lienig & Zeller, 1846, *A. chalcochrysa* Meyrick, 1913, *A. submontana* Frey, 1870.

## Discussion

This study represents preliminary work to describe the diversity of the genus *Argyresthia* in the Neotropics and describes the first species of this family in Central America. All species described are from moderate- to high- elevation areas 1549 m to 2439 m in elevation. Due to the lack of data regarding *Argyresthia* in the Neotropics and the cooler habitat these specimens were collected in, it was necessary to ensure these specimens indeed represented new species and were not part of a range extension for Nearctic conspecifics or were agricultural invasions. Evidence from the phylogenetic inferences using COI supports the monophyly of Argyresthiidae, which has been demonstrated with fewer taxa in past work (Sohn et al. 2013). Both the IQ-TREE maximum likelihood and Bayesian phylogenetic inferences have multiple regions of uncertainty due to soft polytomies representing the inability of the well-conserved COI to resolve all internal relationships in the group. While the topology appears more branching in nature in the IQ-TREE inference, the support is generally low and does not provide additional insight where the Bayesian inference recovered polytomies. Despite this, most species for which there were more than one exemplar were recovered as monophyletic.

Of particular interest is the paraphyly of both *Argyresthia calliphanes* and *A. goedartella*. Meyrick (1913), in the original description, characterized *A. calliphanes* as having a white thorax and *Argyresthia goedartella* a golden one. Forbes (1923) in his Lepidoptera of New York series seconds Meyrick's notes on distinguishing the two species saying of *A. calliphanes*: "Head white, thorax white, wing markings exactly like those of *A. goedartella*", and of *A. goedartella* "thorax bright golden, fore wing white and coppery golden". He also states the range for *A. goedartella* to be "Europe; reported from some places in the United States, in part, at least, in error for *A. calliphanes*". Furthermore, Lewis & Sohn (2015) also note: "This species has been confused with *A. goedartella* L. It is still unclear if both *A. calliphanes* and *A. goedartella* are present in North America. Therefore, some Nearctic records of *A. goedartella* may be due to confusion with *A. calliphanes*. A careful revision of these records is necessary." Pohl et al. (2018) only considered individuals as *Argyresthia calliphanes* if from the type locality of Toronto, Ontario, Canada, but noted concerns by Forbes (1923) and Covell (1984). There are currently no Nearctic specimens in the Cornell University Insect Collection identified as either with a gold/yellowish thorax, which at face value would indicate Forbes' supposition as true. Some examples photographed from Western North America appear to have a gold thorax, but this is difficult to evaluate given cuticle wear often presents itself in images as gold-appearing thoraces. If they do indeed have a gold thorax, this may indicate a movement into North America potentially associated with Beringia. Based on existing taxonomic works, it is prudent to refer to North American specimens identified as *Argyresthia goedartella* as *A. calliphanes* until further treatment of the group which may represent a species complex, as at least three lineages were identified in the phylogenetic inference. Resolving this

taxonomic issue will require extensive sampling across the Nearctic but focused on northeastern North America, southwestern North America, and northwestern North America.

*Argyresthia cupressella* was rendered paraphyletic in both inferences, with one clade, consisting entirely of specimens collected in the District of Columbia, United States of America, coming out sister to *A. glaucinella*, and a clade consisting entirely of specimens from British Columbia, Canada coming out as sister to *A. canadensis* Freeman 1972 + (*A. aureoargentella* Brower, 1953 + *A. deletella*). Considering *A. cupressella* is a western species (type locality Los Angeles, California), and that the representatives from BOLD listed as *A. cupressella* do not resemble *A. cupressella*, this likely represents a misidentification and should not call the species into question. Since most sequences were downloaded from either GenBank or BOLD, and were thus not examined, it is unclear for most species whether species recovered as paraphyletic were due to misidentifications, poor molecular resolution, or indeed represent paraphyletic lineages. Further analysis using a broader Sanger dataset, or ideally, a high-throughput sequencing approach, is preferred to elucidate the relationships of this family more clearly.

The stochastic character mapping suggests that the evolution of this family is closely tied to host plant associations, particularly a grade from angiosperm feeding *Argyresthia* to *Argyresthia* that feed on gymnosperms. Only two species in the clade of primarily Cupressaceae-feeding *Argyresthia* had a reversal to angiosperms—*A. subreticulata* (Sapindaceae) and *A. glaucinella* (multiple angiosperm hosts). No reversals to a gymnosperm host were observed in the angiosperm clade. While the stochastic character mapping should be viewed with some caution due to the lack of resolution provided by a single-locus phylogeny, the consistency of these traits suggests that better resolution will support these initial findings. The ancestral state for the family is hypothesized to potentially Betulaceae-feeding. The ancestral state for the primarily angiosperm clade is Rosaceae-feeding, but there are angiosperm-feeders ancestrally Betulaceae-feeding that grade into a primarily Cupressaceae-feeding clade. While this is a preliminary investigation into host plant associations in these families, the phylogenetic implications of conserved host plant evolutions are notable—there may be a potential to predict host plants for new species or species for which we do not currently have host plant information from with a reasonable degree of confidence. For example, while the newly described species here *A. guatemalae* groups ambiguously among Betulaceae- and gymnosperm-feeding *Argyresthia*, *A. quetzaltenangonella* and *A. iridescentia* group with the clade containing the gymnosperm-feeding *Argyresthia*, *A. quetzaltenangonella* more specifically within a clade ancestrally Cupressaceae-feeding, and *A. iridescentia* within an ancestrally Pinaceae-feeding. The stochastic character mapping likely oversimplifies host plant evolution since the Mk1 model posits equally likely changes for all character states and biologically, it is more likely that host plant switches occur within gymnosperms versus within angiosperms. Similarly, the inclusion of multiple specimens for paraphyletic species also influences the stochastic character mapping yet is also expected to not influence the results broadly.

The following representatives of the subgenus *Blastotere* were included in the phylogenetic inference: *Argyresthia* (*Blastotere*) *amiantella*; *A. (B.) arceuthina*; *A. (B.) bergiella*; *A. (B.) dilectella*; *A. (B.) glabratella*; *A. (B.) illuminatella*; *A. (B.) laevigatella*, *A. (B.) kulfani*, *A. (B.) praecocella*; *A. (B.) svenssoni*; *A. (B.) trifasciata*. These representatives were recovered as paraphyletic, though a few species groups are monophyletic (Figure 6). More phylogenetic and morphological work is necessary to delineate whether the members of this clade that render *Blastotere* paraphyletic are also members of *Blastotere*, meaning *Blastotere* is monophyletic, or if *Blastotere* represents a paraphyletic lineage.

The species descriptions of *Argyresthia* in this paper are by no means exhaustive but rather represent a relatively high degree of diversity returned from a small sample size. There is no doubt that additional sampling in Guatemala, as well as other locales in Central and South America, will bring more species of *Argyresthia* to light.

## Acknowledgements

The authors would like to thank Universidad del Valle de Guatemala for support, especially Jack Schuster and the Centro de Estudios Ambientales y Biodiversidad (CEAB), as well as Tim McCabe

(New York State Museum) for assistance with fieldwork. The authors are also indebted to the Consejo Nacional de Áreas Protegidas (CONAP) for facilitating research, collecting and export permits. This work was supported in part by the Jane E. Brody Undergraduate Research Award, the Fredric N. Gabler '93 Memorial Research Endowment, and the CALS Alumni Association Academic Enrichment Program.

## References

- Alford, D. V. (1978). The true identity of the cherry fruit moth in Britain. *Plant Pathology*, 27(1), 35-37.
- Alford, D. V. (2007). *Pests of Fruit Crops, A Color Handbook*. Manson Publishing Ltd.
- Ahlberg, O. (1927). Ronnbarsmalen, *Argyresthia conjugella* Zell. En redogorelse for undersokningar aren 1921-1926. *Meddel Nr 324 fran Centralanstalten for forsoksvasendet pa jordbruksomradet*. Lantbruksentomologiska avdelningen.
- Belosel's Kaya, Z. C. (1963). The mountain-ash moth *Argyresthia conjugella* Zell. (Lep. Yponomeutidae) as a pest of the fruits of apples and mountain-ash. *Entomologicheskoye Obozreniye*, 4, 720-729.
- Bengtsson, B. Å., & Johansson, R. (2011). Review of the unicolorous species of the subgenus *Blastotere* (Lepidoptera, Argyresthiidae) with descriptions of *Argyresthia svenssoni* sp. nov. and *A. kulfani* sp. nov. *Entomologisk Tidskrift*, 132, 257-274.
- Boyko, J. D. & Beaulieu, J. M. (2021). Generalized hidden Markov models for phylogenetic comparative datasets. *Methods in Ecology and Evolution*, 12, 468-478.
- Braun, A. F. (1910). New species of Tineina from California. *Entomological News*, 21, 171-179.
- Braun, A. F. (1940). Notes and new species in the Yponomeutoid Group (Microlepidoptera). *Transactions of the American Entomological Society*, 66(1079), 273-282.
- Brower, A. E. (1953). Three new species of microlepidoptera (Olethreutidae, Glyphipterygidae and Yponomeutidae). *Annals of the Entomological Society of America*, 46(1), 95-98.
- Busck, A. (1907). Revision of the American moths of the genus *Argyresthia*. *Proceedings of the United States National Museum*, 32, 5-24.
- Busck, A. (1915). Descriptions of new North American Microlepidoptera. *Proceedings of the Entomological Society of Washington*, 17(2), 79-94.
- Butler, A. G. (1883). III. Heterocerous Lepidoptera collected in Chile by Thomas Edmonds, Esq. *Transactions of the Entomological Society of London*, 1883, 49-90.
- Carter, D. J. (1984). *Pest Lepidoptera of Europe: with special reference to the British Isles*. Dr W. Junk Publishers.
- Covell Jr., C. V. (1984). *A Field Guide to the moths of Eastern North America*. Houghton Mifflin Co.
- Clerck, C. A. (1759). *Icones Insectorum rariorum. Sectio Prima. Insectorum rariorum cum nominibus eorum trivialibus locisque e C. Linnaei systema naturae allegatis*. Stockholm.
- Csóka, G. (2001). Recent Invasions of Five Species of Leaf mining Lepidoptera in Hungary. In A. M. Liebhold, M. L. Mcmanus, I. S. Otvos & S. L. C. Fosbroke. *Proceedings: Integrated Management and Dynamics of Forest Defoliating Insects* (pp. 31-36). USDA Forest Service, Northeastern Research Station. <https://doi.org/10.2737/NE-GTR-277>
- Folmer, O., Black, M., Hoeh, W., Lutz, R., & Vrijenhoek, R. (1994). DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology*, 3, 294-299.
- Forbes, W. T. M. (1923). *The Lepidoptera of New York and Neighboring States*. Cornell University.
- Freeman, T. N. (1960). Needle-Mining Lepidoptera of Pine in North America. *The Canadian Entomology*, 92(Supplement 16), 1-51.
- Freeman, T. N. (1972). The Coniferous Feeding Species of *Argyresthia* in Canada (Lepidoptera: Yponomeutidae). *The Canadian Entomology*, 104, 687-697.
- Frey, H. (1870). Ein Beitrag zur Kenntnis der Microlepidopteren. *Mittheilungen der Schweizerischen Entomologischen Gesellschaft*, 3(5), 244-256, (6): 277-296.
- Friese, G. (1969). Beiträge zur Insekten-Fauna der DDR: Lepidoptera-Argyresthiidae. *Beiträge zur Entomologie*, 19(7/8), 693-752.
- Hepner, J. B. (2008). Shiny Head-Standing Moths (Lepidoptera: Argyresthiidae). In J. L. Capinera (ed.). *Encyclopedia of Entomology*. Springer.

- Herrich-Schffer, G. A. W. (1855). *Systematische Bearbeitung der Schmetterlinge von Europa, zugleich als Text, Revision und Supplement zu Jakob Hübner's Sammlung europäischer Schmetterlinge* (Vol. 5). G. J. Marz.
- Hoang, D. T., Chernomor, O., Von Haeseler, A., Minh, B. Q., & Vinh, L. S. (2018). UFBBoot2: improving the ultrafast bootstrap approximation. *Molecular Biology and Evolution*, 35, 518-522.
- Hübner, J. (1825). *Verzeichnis bekannter Schmettlinge*. Augsburg.
- Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K., Von Haeseler, A., & Jermini, L. S. (2017). ModelFinder: fast model selection for accurate phylogenetic estimates. *Natural Methods*, 14, 587.
- Katoh, K., & Standley, D. M. (2013). MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability. *Molecular Biology and Evolution*, 30, 772-780.
- Jaastad, G. (2007). Late dormant rapeseed oil treatment against black cherry aphid and cherry fruit moth in sweet cherries. *Journal of Applied Entomology*, 131(4), 284-288.
- Korchagin, V. N. (1988). Pestsof fruit tress. *Zashchita Rasten (Moskva)*, (3), 48-52.
- Lampa, S. (1905). Rönnbärsmalen (*Argyresthia conjugella* Zell.). *Entomologisk Tidskrift*, 26, 27.
- Lewis, J., & Sohn, J.-C. (2015). Lepidoptera: Yponomeutoidea I (Argyresthiidae, Attevidae, Praydidae, Scythropiidae, and Yponomeutidae). *World Catalogue of Insects*, 12. Brill.
- Lienig, F., & Zeller, P. C. (1846). Lepidopterologische Fauna von Livland und Curland. Bearbeitet von Friederike Lienig, geb. Berg, mit Anmerkungen von P. C. Zeller. *Isis von Oken*, 39(3/4), 202-302.
- Linnaeus, C. (1758). *Systema Naturae per Regna Tria Nature, secundum Classes, Ordines, Genera, Species, cum characteribus, differentiis, synonymis, locis. Edition 10*. Laurentii Salvii.
- Liu, T., Wang, S., & Li, H. (2017). Review of the genus *Argyresthia* Hübner, [1825] (Lepidoptera: Yponomeutoidea: Argyresthiidae) from China, with descriptions of forty-three new species. *Zootaxa*, 4292(1), 1-135.
- Looney, C., Lagasa, E., & Passoa, S. (2017). Distribution, biology, and identification of *Argyresthia pruniella* in Washington State. *Journal of the Lepidopterist Society*, 71(2), 117-122.
- Maddison, W. P., & Maddison, D. R. (2018). *Mesquite: a modular system for evolutionary analysis*. Version 3.6. <http://www.mesquiteproject.org>
- Meyrick, E. (1912-1916). *Exotic Microlepidoptera*, 1. Thornhanger.
- Meyrick, E. (1916-1923). *Exotic Microlepidoptera*, 2. Thornhanger.
- Meyrick, E. (1930-1936). *Exotic Microlepidoptera*, 4. Thornhanger.
- Minh, B. Q., Nguyen, M. A. T., & Von Haeseler, A. (2013). Ultrafast approximation for phylogenetic bootstrap. *Molecular Biology and Evolution*, 30(5), 1188-1195.
- Moriuti, S. (1974). A new Formosan Argyresthiidae (Lepidoptera). *Transactions of the Lepidopterological Society of Japan*, 25, 113-114.
- Moriuti, S. (1977). *Fauna Japonica, Yponomeutidae s. lat. (Insecta: Lepidoptera)*. Keigaku Publishing Company.
- Nguyen, L.-T., Schmidt, H. A., Von Haeseler, A., & Minh, B. Q. (2015). IQ-TREE: A fast and effective stochastic algorithm for estimating maximum likelihood phylogenies. *Molecular Biology and Evolution*, 32, 268-274.
- Packard, A. S. (1871). *First annual report on the injurious and beneficial insects of Massachusetts*. Wright & Potter.
- Pennell, M. W., Eastman, J. M., Slater, G. J., Brown, J. W., Uyeda, J. C., Fitzjohn, R. G., Alfaro, M. E., & Harmon, L. J. (2014). Geiger v2. 0: an expanded suite of methods for fitting macroevolutionary models to phylogenetic trees. *Bioinformatics*, 30, 2216-2218.
- Pohl, G. R., Landry, J. F., Schmidt, B. C., Lafontaine, J. D., Troubridge, J. T., Macaulay, A. D., Nieukerken, E. J., Van, Dewaard, J. R., Dombroskie, J. J., Klymko, J., & Nazari, V. (2018). *Annotated checklist of the moths and butterflies (Lepidoptera) of Canada and Alaska*. Pensoft.
- Ratzeburg, J. T. C. (1840). *Die Forst-insekten oder Abbildung und Beschreibung der in der Wäldern Preusens und der Nachbarstaaten als schädlich oder nützlich bekannt geworden Insekten; In systematischer Folge und mit besonderer Rücksicht auf die Vertilgung der Schädlichen. Zweiter Theil. Die Falter* (Vol. 2). Nicolaische Buschhandlung.
- Rebel, H. (1940). Die Lepidopterenfauna des Azorischen Archipels. *Societas Scientiarum Fennica. Commentationes Biologicae*, 8(1), 35-60.
- Revell, L. J. (2012). phytools: an R package for phylogenetic comparative biology (and other things). *Methods in Ecology and Evolution*, 2, 217-223.
- Rstudio Team (2022). *RStudio: Integrated Development for R*. RStudio, Inc., Boston. <http://www.rstudio.com/>
- Schøyen, T. H. (1913). *Beretning om Skadeinsekter og Plantesygdomme*. Kristiania.
- Sharma, J. P., Khajuria, D. R., & Dogra, G. S. (1988). Studies on the apple fruit moth, *Argyresthia conjugella* Zeller (Yponomeutidae: Lepidoptera): Identification, distribution, and extent of damage in India. *International Journal of Pest Management*, 34(2), 189-192.

- Shorthouse, D. P. (2010). *SimpleMappr*, an online tool to produce publication-quality point maps. <http://www.simplemappr.net>
- Shreyner, Y. F. (1906). Principal insects attacking fruit crops, particularly in the north and methods for exterminating them. *Plodovostova*, 23, 55.
- Sohn, J.-C., Regier, J. C., Mitter, C., Davis, D., Landry, J.-F., Zwick, A., & Cummings, M. P. (2013). A Molecular Phylogeny for Yponomeutoidea (Insecta, Lepidoptera, Ditrysia) and Its Implications for Classification, Biogeography, and the Evolution of Host Plant Use. *PLoS One*, 8(1), e55066. <https://doi.org/10.1371/journal.pone.0055066>
- Stapley, J. H. (1934). The apple fruit miner (*Argyresthia conjugella* Zell.). *Journal of South-Eastern Agricultural College*, 34, 87, 92.
- Staudinger, O. (1871). Macrolepidoptera. In O. Staudiner & M. Wocke. *Catalog der Lepidopteren des Europäischen Faunengebiets* (pp. 473). Hermann Burdach.
- Stritch, L. (2018). *Castanea dentata*. The IUCN Red List of Threatened Species.
- Tenow, O., Nilssen, A. C., Holmgren, B., & Elverum, F. (1999). An insect (*Argyresthia retinella*, Lep., Yponomeuti-dae) outbreak in northern birch forests, released by climatic changes? *The Journal of Applied Ecology*, 36(1), 111-122.
- Tiborne, G., & Kálmán, S. (2000). Appearance of *Argyresthia trifasciata* Staudinger, 1871 (Lepidoptera, Yponomeutidae) on *Juniperus* in Hungary. *Növényvédelem*, 36, 301-304.
- Turner, A. J. (1913). Studies in Australian Microlepidoptera. *Proceedings of the Linnean Society of New South Wales*, 38, 174-228.
- Walsingham, L. (1890). Notes on the genus *Argyresthia* Hb., with descriptions of new species. *Insect Life*, 3(3), 118.
- Wimshurst, F. M. (1928). Preliminary note on the cherry fruit moth (*Argyresthia nitidella* Fabr). *Journal of South-Eastern Agricultural College*, 25, 86-89.
- World Conservation Monitoring Centre (1996). *Argyresthia castaneella*. *The IUCN Red List of Threatened Species 1996*, e.T2100A9242798.
- Zeller, P. C. (1839). Versuch einer naturgemässen Eintheilung der Schaben. *Isis von Oken*, 32(3), 167-220.
- Zeller, P. C. (1847). Die *Argyresthien*. *Linnaea entomologica*, 2, 234-302.
- Zeller, P. C. (1873). Beiträge zur Kenntniss der nordamerikanischen Nachtfalter, besonders der Microlepidopteren. *Verhandlungen der kaiserlich-königlichen Zoologische-botanischen Gesellschaft in Wien*, 23(3), 201-334.
- Zeller, P. C. (1877). Exotische Microlepidoptera. *Horae Societatis Entomologicae Rossicae*, 13, 3-498.

\*Jacob A. Gorneau  
Department of Entomology  
Cornell University  
Ithaca, NY 14853  
EE.UU. / USA  
E-mail: [jgorneau@calacademy.org](mailto:jgorneau@calacademy.org)  
<https://orcid.org/0000-0001-9230-9774>

y/ and

Institute for Biodiversity Science and Sustainability  
California Academy of Sciences  
55 Music Concourse Drive  
San Francisco, CA, 94118  
EE.UU. / USA

Loren D. Jones  
Department of Entomology  
Cornell University  
Ithaca, NY 14853  
EE.UU. / USA  
E-mail: [ldj37@cornell.edu](mailto:ldj37@cornell.edu)  
<https://orcid.org/0000-0001-8348-8416>

José Monzón-Sierra  
Centro de Estudios Ambientales y Biodiversidad  
Universidad del Valle de Guatemala  
18 Avenida 11-95 Zona 15, Vista Hermosa III  
Guatemala, C. A.  
GUATEMALA / GUATEMALA  
E-mail: [jmonzon@uvg.edu.gt](mailto:jmonzon@uvg.edu.gt)  
<https://orcid.org/0000-0002-0733-1895>

Jason J. Dombroskie  
Department of Entomology  
Cornell University  
Ithaca, NY, 14853  
EE.UU. / USA  
E-mail: [jjd278@cornell.edu](mailto:jjd278@cornell.edu)  
<https://orcid.org/0000-0002-7232-6507>


\*Autor para la correspondencia / *Corresponding author*

(Recibido para publicación / *Received for publication* 22-VII-2022)


(Revisado y aceptado / *Revised and accepted* 20-IX-2022)

(Publicado / *Published* 30-III-2023)


**Derechos de autor:** El autor(es). Este es un artículo de acceso abierto distribuido bajo los términos de la Licencia de Reconocimiento 4.0 Internacional de Creative Commons (CC BY 4.0), que permite el uso, distribución y reproducción sin restricciones en cualquier medio, siempre que se cite al autor original y la fuente. / **Copyright:** The author(s). This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



**Figure 1.** *Argyresthia quetzaltenangonella* Gorneau & Dombroskie, sp. nov., holotype, ♂. A. Adult. B. Coremata of abdomen. C. Male genitalia with phallus removed. D. phallus.




**Figures 2-3.** 2. Dorsal habitus (A), ductus bursae and corpus bursae (B); and genital capsule (C) of holotype *Argyresthia guatemala* Gorneau & Dombroskie, sp. nov. 3. Genital capsule, ductus bursae (A); abdomen (B); and corpus bursae (C) of slide of dissected female *Argyresthia deletella* (USNM 76603).



**Figures 4-5.** 4. Dorsal habitus (A), phallus (B), and genital capsule with valves spread (C) of holotype *Argyresthia iridescentia* Gorneau & Dombroskie, sp. nov. 5. Maps with holotype localities of new *Argyresthia* species made using SimpleMappr (SHORTHOUSE 2010). Holotype locality of *Argyresthia iridescentia* Gorneau & Dombroskie, sp. nov. in black square, of *A. guatemala* Gorneau & Dombroskie, sp. nov. in black star, and of *A. quetzaltenangonella* Gorneau & Dombroskie, sp. nov. in white circle. Paratype locality of *A. guatemala* Gorneau & Dombroskie, sp. nov. in white star.










Figure S2. Phylogenetic plot showing differences between IQ-TREE and MrBayes phylogenetic inferences.

## Supplementary Material

Table SI. Sequences downloaded from GenBank and BOLD for analysis.

Taxon	GenBank Accession(s)	BOLD Voucher(s)
<i>Argyresthia abdominalis</i> Zeller, 1839	HM386989, HM871642,	-
<i>Argyresthia abies</i> Freeman, 1972	GU095507, MG358614	-
<i>Argyresthia abies</i> Freeman, 1972	MG358614	-
<i>Argyresthia affinis</i> Braun, 1940	-	LNAUS5768-14, LNAUS5767-14, LNAUS5766-14
<i>Argyresthia albistria</i> (Haworth, 1828)	HM391874, HM875669	-
<i>Argyresthia alternatella</i> Kearfott, 1908	GU800182, GU800969	-
<i>Argyresthia amiantella</i> (Zeller, 1847)	JF859683, JF859684	-
<i>Argyresthia amnettella</i> Busck, 1907	KM541487, KM549165	-
<i>Argyresthia arceuthina</i> Zeller, 1839	HM876038, HM872265	-
<i>Argyresthia assimilis</i> Moriuti, 1977	LC421504, LC421505	-
<i>Argyresthia atlanticella</i> Rebel, 1940	MT072686, MT072687	-
<i>Argyresthia aureoargentella</i> Brower, 1953	MG361900, MG362496	-
<i>Argyresthia aurulentella</i> Stainton, 1849	HM876198, HM871641	-
<i>Argyresthia austerella</i> Zeller, 1873	HQ989655, KF491556	-
<i>Argyresthia bergiella</i> (Ratzeburg, 1840)	HM874190, HM874453	-
<i>Argyresthia bonnetella</i> (Linnaeus, 1758)	HM873448, HM873449	-
<i>Argyresthia brockeella</i> (Hübner, [1813])	GU707321, HM873792	-
<i>Argyresthia calliphanes</i> Meyrick, 1913	-	LNAUS5784-14, LNAUS5783-14, LNAUS5782-14, LNAUS5781-14, LNAUS5778-14
<i>Argyresthia canadensis</i> Freeman, 1972	GU095510, GU095511	-
<i>Argyresthia chalcocrysa</i> Meyrick, 1918	-	LNAUS5786-14
<i>Argyresthia conjugella</i> Zeller, 1839	GU690418, HM391873	-
<i>Argyresthia cupressella</i> Walsingham, 1891	-	MNAD219-07, MNAD218-07, LNAUT2196-14, LNAUT2190-14, LNAUS5788-14, DUNLP350-08, DUNLP349-08, DUNLP348-08
<i>Argyresthia curvella</i> (Linnaeus, 1764)	GU707322	-
<i>Argyresthia curvella</i> (Linnaeus, 1764)	HM875765	-
<i>Argyresthia deletella</i> Zeller, 1873	-	LNAUS5789-14
<i>Argyresthia dilectella</i> Zeller, 1847	HM875177, HM876037	-
<i>Argyresthia flexilis</i> Freeman, 1960	KT136150, KT139486	-
<i>Argyresthia freyella</i> Walsingham, 1891	-	MNAA065-07, MNAA064-07, MNAA063-07, MNAA062-07, MNAA061-07
<i>Argyresthia fundella</i> Fischer von Röslerstamm, 1834	GU706913, KP253578	-
<i>Argyresthia glabratella</i> (Zeller, 1847)	GU707214, HM875937	-
<i>Argyresthia glaucinella</i> Zeller, 1839	GU707319, HM386995	-
<i>Argyresthia goedartella</i> (Linnaeus, 1758)	GU095513, GU096217, KX043548, KX044037	-
<i>Argyresthia illuminatella</i> Zeller, 1839	KP150223, KP253410	-
<i>Argyresthia ivella</i> (Haworth, 1828)	-	LON3240-16
<i>Argyresthia kulfani</i> Bengtsson & Johansson, 2012	HQ955402	-
<i>Argyresthia laevigatella</i> (Herrich-Schäffer, 1855)	HM876340, KQ955511	-

<i>Argyresthia laricella</i> Kearfott, 1908	MG499568	-
<i>Argyresthia libocedrella</i> Busck, 1916	-	LNAUS5809-14
<i>Argyresthia notoleuca</i> (Turner, 1913)	HQ922352, HQ922353	-
<i>Argyresthia oreasella</i> Clemens, 1861	GU095514, GU096219	-
<i>Argyresthia praecocella</i> Zeller, 1839	KT782576	-
<i>Argyresthia pruniella</i> (Clerck, 1759)	FJ412159, FJ412160	-
<i>Argyresthia pulchella</i> Lienig & Zeller, 1846	HM875282, HM875758	-
<i>Argyresthia pygmaeella</i> ([Denis & Schiffermüller], 1775)	GU679758, HM865123	-
<i>Argyresthia quadristrigella</i> Zeller, 1873	GU801642, KT130002	-
<i>Argyresthia reticulata</i> Staudinger, 1877	-	LON6890-18
<i>Argyresthia retinella</i> Zeller, 1839	HM875671, HQ570278	-
<i>Argyresthia ruidosa</i> Braun, 1940	-	LNAUT3197-14
<i>Argyresthia semifusca</i> (Haworth, 1828)	HM387223	-
<i>Argyresthia semitestacella</i> (Curtis, 1833)	HQ955406, HQ955519	-
<i>Argyresthia sorbiella</i> (Treitschke, 1833)	HM875670, HM875977	-
<i>Argyresthia spinosella</i> Stainton, 1849	HM873477, HM873479	-
<i>Argyresthia submontana</i> Frey, [1871]	-	-
<i>Argyresthia subreticulata</i> Wlsingham, 1882	GU088278, GU801315	-
<i>Argyresthia svenssoni</i> Bengtsson & Johansson, 2011	HM873235, JF853437	-
<i>Argyresthia thuiella</i> (Packard, 1871)	GU095515, GU095516	-
<i>Argyresthia trifasciae</i> Braun, 1910	-	LNAUS5832-14, LNAUS5833-14
<i>Argyresthia trifasciata</i> Staudinger, 1871	GU706912, GU707204	-
<i>Argyresthia tsuga</i> Freeman, 1972	MG502448	-
<i>Argyresthia</i> sp. 1 TL-2017	KY570298, KY570300	-
<i>Argyresthia</i> sp. 2 TL-2017	KY570301, KY570302	-
<i>Embola ionis</i> (Clarke, 1952)	JF841791	-
<i>Lyonetia candida</i> Braun, 1916	KF808908	-
<i>Ypsolopha dentella</i> (Fabricius, 1775)	GU0914191	-
<i>Acrolepiopsis assectella</i> (Zeller, 1839)	EF380034	-
<i>Atteva aurea</i> (Fitch, 1856)	GU013569	-
<i>Bedellia somnulentella</i> (Zeller, 1847)	GU828759	-
<i>Plutella xylostella</i> (Linnaeus, 1758)	DQ076339	-
<i>Yponomeuta malinellus</i> Zeller, 1838	FJ413018	-
<i>Prays fraxinella</i> (Bjerkander, 1784)	GU706347	-

**Table SII.** Matrix used in ancestral state reconstruction with host information from Lewis and Sohn (2015) reduced to family level or listed as multiple angiosperm families if species feeds on multiple families. Multiple representatives retained from original phylogenetic analysis for specimens that were not recovered as monophyletic. Accession information kept for efficient retrieval of specific specimen information.

Taxon	Accession Number(s)	Host Family
<i>Argyresthia abdominalis</i> Zeller, 1839	HM386989	Cupressaceae
<i>Argyresthia abies</i> Freeman, 1972	GU095507, MG358614	Pinaceae
<i>Argyresthia affinis</i> Braun, 1940	LNAUS5767-14	Cupressaceae
<i>Argyresthia albistria</i> (Haworth, 1828)	HM391874	Rosaceae
<i>Argyresthia alternatella</i> Kearfott, 1908	GU800182	Cupressaceae
<i>Argyresthia annettella</i> Busck, 1907	KM541487	Cupressaceae
<i>Argyresthia arceuthina</i> Zeller, 1839	HM876038	Cupressaceae

<i>Argyresthia aureoargentella</i> Brower, 1953	MG361900	Cupressaceae
<i>Argyresthia aurulentella</i> Stainton, 1849	HM876198	Cupressaceae
<i>Argyresthia bonnetella</i> (Linnaeus, 1758)	HM873448	Rosaceae
<i>Argyresthia brockeella</i> (Hübner, [1823])	GU707321	Betulaceae
<i>Argyresthia calliphanes</i> Meyrick, 1913	LNAUS5782-14	Betulaceae
<i>Argyresthia canadensis</i> Freeman, 1972	GU095510	Cupressaceae
<i>Argyresthia conjugella</i> Zeller, 1839	GU690418	Rosaceae
<i>Argyresthia cupressella</i> Walsingham, 1891	LNAUS5788-14, MNAD219-07	Cupressaceae
<i>Argyresthia curvella</i> (Linnaeus, 1761)	GU707322	Multiple Angiosperm Families
<i>Argyresthia dilectella</i> Zeller, 1839	HM875177	Cupressaceae
<i>Argyresthia flexilis</i> Freeman, 1960	KT136150	Pinaceae
<i>Argyresthia freyella</i> Walsingham, 1891	MNAA062-07	Cupressaceae
<i>Argyresthia fundella</i> Fischer von Röslerstamm, 1834	GU706913	Pinaceae
<i>Argyresthia glabratella</i> Zeller, 1847	HM875937	Pinaceae
<i>Argyresthia glaucinella</i> Zeller, 1839	GU707319	Multiple Angiosperm Families
<i>Argyresthia goedartella</i> (Linnaeus, 1758)	KX044037	Multiple Angiosperm Families
<i>Argyresthia illuminatella</i> Zeller, 1839	KP150223	Pinaceae
<i>Argyresthia ivella</i> (Haworth, 1828)	LON3240-16	Multiple Angiosperm Families
<i>Argyresthia laevigatella</i> (Herrich-Schäffer, 1855)	HM876340	Pinaceae
<i>Argyresthia loricella</i> Kearfott, 1908	MG499568	Pinaceae
<i>Argyresthia libocedrella</i> Busck, 1916	LNAUS5809-14	Cupressaceae
<i>Argyresthia oreasella</i> Clemens, 1861	GU095514	Rosaceae
<i>Argyresthia praecocella</i> Zeller, 1839	KT782576	Cupressaceae
<i>Argyresthia pruniella</i> (Clerck, 1759)	FJ412159	Rosaceae
<i>Argyresthia pygmaeella</i> ([Denis & Schiffmüller], 1775)	GU679758	Salicaceae
<i>Argyresthia quadririgella</i> Zeller, 1873	KT130002	Cupressaceae
<i>Argyresthia reticulata</i> Staudinger, 1877	LON6890-18	Cupressaceae
<i>Argyresthia retinella</i> Zeller, 1839	HM875671	Betulaceae
<i>Argyresthia semifusca</i> (Haworth, 1828)	HM387223	Rosaceae
<i>Argyresthia semitestacella</i> (Curtis, 1833)	HQ955406	Fagaceae
<i>Argyresthia sorbiella</i> (Treitschke, 1833)	HM875670	Rosaceae
<i>Argyresthia spinosella</i> Stainton, 1849	HM873477	Sapindaceae
<i>Argyresthia subreticulata</i> Walsingham, 1882	GU801315	Cupressaceae
<i>Argyresthia thuiella</i> (Packard, 1871)	GU095515	Cupressaceae
<i>Argyresthia trifasciae</i> Braun, 1910	LNAUS5832-14, LNAUS5833-14	Cupressaceae
<i>Argyresthia trifasciata</i> Staudinger, 1871	GU706912	Cupressaceae
<i>Argyresthia tsuga</i> Freeman, 1972	MG502448	Pinaceae